

Increasing temperature and pH can facilitate reductions of cephapirin and antibiotic resistance genes in dairy manure slurries

Article

Accepted Version

Li, M. M., Ray, P., Teets, C., Pruden, A., Xia, K. and Knowlton, K. F. (2020) Increasing temperature and pH can facilitate reductions of cephapirin and antibiotic resistance genes in dairy manure slurries. *Journal of Dairy Science*, 103 (3). pp. 2877-2882. ISSN 0022-0302 doi: <https://doi.org/10.3168/jds.2019-17453> Available at <https://centaur.reading.ac.uk/87405/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.3168/jds.2019-17453>

Publisher: American Dairy Science Association

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

1 INTERPRETIVE SUMMARY: *Li et al* (2019). “*Short communication: Raising temperature and*
2 *pH can facilitate reductions of cephapirin and antibiotic resistance genes in dairy manure*
3 *slurries*”. Quantifying the fate of antibiotics and antibiotic resistance genes (ARGs) exposed to
4 various temperatures and pH adjustments in manure slurries will help guide manure management
5 practices to minimize the emergence and subsequent spreading of antimicrobial resistance. In the
6 current study, effects of temperature and initial pH shock on cephapirin and ARGs in dairy
7 manure slurries were investigated using a microcosm setup. Our results suggest a change in
8 temperature or initial pH adjustment during storage of dairy manure slurries could mitigate the
9 spread of resistance.

Short communication: Raising temperature and pH can facilitate reductions of cephalosporin and antibiotic resistance genes in dairy manure slurries

Meng M. Li^{†1}, Partha Ray[§], Christy Teets[†], Amy Pruden[‡], Kang Xia*, Katharine F. Knowlton[†]

[†]Department of Dairy Science, [‡]Department of Civil and Environmental Engineering, and

*Department of Crop and Soil Environmental Sciences, Virginia Tech, Blacksburg, VA, USA;

[§]Department of Animal Sciences, School of Agriculture, Policy and Development, University of

Reading, Reading RG6 6AR, UK;

¹ Corresponding author (mengli@vt.edu)

Abstract

Quantifying antibiotics and antibiotic resistance genes (ARGs) in manure exposed to various temperature and pH treatments could guide the development of cost-effective manure handling methods to minimize the spread of antibiotic resistance following land application of manure. This study aimed to investigate the effect of various temperatures and initial pH shocks on the persistence of a cephalosporin antibiotic and ARGs in dairy manure slurries. Feces and urine were collected from five healthy dairy cows administered with cephapirin (cephalosporin antibiotic) at dry-off via intramammary infusion, and mixed with sterile water to generate manure slurries. In a 28-day incubation study, dairy manure slurries were either continuously exposed to one of three temperatures (10, 35, and 55°C) or received various initial pH (5, 7, 9, and 12) shocks. Cephapirin was detected in the initial samples and on day 1 following all treatments, but it was undetectable thereafter. This indicates that cephapirin can be rapidly degraded irrespective of temperature and pH treatments. However, degradation was greater on day 1 with the mesophilic (35°C) and thermophilic (55°C) environments compared to the psychrophilic environment at 10°C ($P < 0.001$). Increasing pH beyond neutral also accelerated degradation as cephapirin concentrations were lower on day 1 after initial alkaline adjustments (pH 9 and 12) than neutral and acidic adjustments (pH 7 and 5; $P < 0.001$). No significant effect of temperature or initial pH was observed on abundances of a beta-lactam ARG, *cfxA*, and a tetracycline ARG, *tet(W)*, implying that bacteria that encoded *cfxA* or *tet(W)* genes were not sensitive to temperature or pH in dairy manure slurries. However, abundances of a macrolide ARG, *mefA*, were decreased in the psychrophilic and thermophilic environments, and also following exposure to a strong alkaline shock (pH 12). Our results suggest that increasing temperature or pH during storage of dairy manure slurries could be used together with other on-

farm practices that are tailored to reduce the transfer of ARGs from manure to the environment following land application.

Key words: antibiotic resistance genes; cephalosporin; dairy manure slurry

Short Communication

In 2017, approximately 10.93 million kg of antimicrobial drugs were sold in the United States for use in food-producing animals, and ~50% of that was used in cattle (USFDA, 2018). Previous studies indicated that up to 90% of administered antibiotics are eliminated from animals' bodies through feces or urine (Kemper, 2008; Ray et al., 2014), implying that manure generated from animal production represents a major route of antibiotics transfer to the environment. The presence of antibiotics, even at a very low concentrations, can contribute to emergence of antibiotic resistance genes (ARGs) and selection of antibiotic resistant bacteria (Knapp et al., 2008; Martínez, 2008; Sandegren, 2014). Many studies have demonstrated that land application of untreated, antibiotic-laden manure can substantially enhance the abundance of ARGs in agricultural soils (Fahrenfeld et al., 2014; Hu et al., 2016; Gou et al., 2018). These ARGs can be transferred to animals and humans through drinking water or the food chain, leading to decreased effectiveness of subsequent antibiotic therapies (Alcaine et al., 2005; Tello et al., 2012). Therefore, it is vital to develop cost-effective methods to degrade antibiotics in manure before land application to mitigate the dissemination of antibiotic resistance.

Manure management practices such as composting and anaerobic digestion could effectively increase degradation of antibiotics and reduce the prevalence of ARGs (Qian et al., 2016; Sun et al., 2016; Ray et al., 2017; Gou et al., 2018). Ray et al. (2017) indicated that thermophilic temperature during the manure composting process could completely degrade cephalosporin, lincosamide, and sulfonamide antibiotics. Sun et al. (2016) reported that the

abundances of 8 of 10 detected ARGs declined during thermophilic anaerobic digestion of dairy manure. However, high maintenance and operation costs, as well as high technical requirements, limit implementation of composting or anaerobic digester operations on many farms (Pandyaswargo and Premakumara, 2014; Garfí et al., 2016). Physico-chemical factors, such as temperature, pH, CO₂ and O₂ levels, likely explain the effect of composting and anaerobic digestion on persistence of antibiotics and ARGs (Sun et al., 2016; Youngquist et al., 2016; Li et al. 2019). The objective of the current study was to isolate two of these factors, temperature and pH, and evaluate their independent effects on antibiotic and ARG persistence. Understanding these mechanisms will inform future development of best management practices for preventing the spread of antibiotic resistance from livestock farms.

Five dairy cows at the end of lactation with similar body weight (574.9 ± 32.8 kg) were fitted with urinary catheters to allow separate collection of feces and urine and moved to individual metabolic stalls at the Virginia Tech Dairy Center (Blacksburg, VA). After 24 h of acclimation to the barn and catheters, the cows were treated with a single dose of 300 mg cephapirin benzathine (Boehringer Ingelheim Vetmedica, Inc., St. Joseph, MO) into each of the four mammary glands through the teat as in the standard dry-off protocol (dry cow antibiotic treatment 60 days before calving) of the Virginia Tech Dairy Center. All cows were allowed ad libitum access to water and a ration formulated to meet their nutritional needs.

Feces and urine were collected from day 1 (the day cephapirin was administered) through day 3 following treatments, and then all feces and urine excreted by a cow over the three days were combined to form one homogenous pool of manure. Approximately 1 kg of the manure mixture was mixed with sterile water to generate a manure slurry with a final solid content of 5%. Aliquots (200 mL) of slurry were transferred into glass beakers (400 mL) and incubated at

10, 35, and 55°C or at 25°C with initial pH adjustments to 5, 7, 9, or 12 using 1 M (mol/L) HCl (for pH 5 and 7) or NaOH (for pH 9 and 12) solution. Each treatment had 4 replicates. Beakers were covered with aluminum foil with a hole in the middle to maintain aerobic conditions. Beakers were weighed daily, and sterile water was added to each beaker to replace weight loss due to water evaporation. Samples were collected on days 0, 1, 3, 7, and 28 and were immediately stored at -20°C for future analysis.

Cephapirin concentrations were quantified following the method developed by Ray et al. (2014). To quantify ARGs, frozen manure slurries were freeze-dried, and then DNA was extracted from the freeze-dried manure samples using the QIAamp DNA stool extraction kit (Qiagen, Valencia, CA) following the manufacturer's instructions. Quantitative PCR (qPCR) was performed to quantify a beta-lactam ARG, *cfxA*, a macrolide ARG, *mefA*, and a tetracycline ARG, *tet(W)*, using an EvaGreen assay (Biotium, Hayward, CA) with previously reported primers (Aminov et al., 2001; S6ki et al., 2011; Looft et al., 2012). 16S rRNA gene copy numbers were determined using the method by Suzuki et al. (2000). Bacterial gene abundances were represented as 16S rRNA gene copies per g dry manure, and ARG relative abundances were expressed as the proportion to 16S rRNA gene abundance.

Data analyses were conducted in R software (version 3.3.0; R development Core Team, 2015). All qPCR results were normalized using log base 10 transformation. The temperature or initial pH effect on cephalosporin concentrations and ARGs was analyzed in a mixed statistical model with temperature or initial pH as a fixed effect and incubation day as a repeated measurement using the gls function in the nlme package (Pinheiro et al., 2018). The first-order autoregressive covariance with heterogeneous variance of time were considered in the repeated

measures ANOVA model. The Tukey's honestly significant difference post hoc test was used to test differences among treatments, and $P < 0.05$ was considered significant.

Cephapirin was present in manure slurries on day 0 (initial samples) and day 1 following either temperature or initial pH adjustment, but it was undetectable thereafter. On day 0, the mean concentration of cephapirin was 12.5 ± 1.3 ng per g dry manure; this rapidly decreased to less than 1 ng per g dry manure on day 1 regardless of temperature or pH treatments (Figure 1). Ray et al. (2017) reported that cephapirin was only present in day 0 compost samples. Gilbertson et al. (1990) indicated that another cephalosporin (ceftiofur) was quickly degraded within 8 h in cattle feces. Thus, our result was consistent with the previous studies, suggesting that cephapirin has a poor stability in dairy manure slurries. Compared to the psychrophilic environment (10°C), the mesophilic (35°C) and thermophilic (55°C) environments decreased cephapirin concentrations on day 1, with no difference between the mesophilic and thermophilic conditions (Figure 1A).

Cephapirin belongs to the β -lactam class of antibiotics which have a basic β -lactam ring structure for the antibacterial activity. Ray et al. (2017) indicated that cephapirin was unstable at high temperature and Cha et al. (2006) demonstrated that the β -lactam ring has poor stability in animal manure and lagoon effluent. Wagner et al. (2011) observed that multiple β -lactamases were detected in bacteria isolated from the digestive tract of cows. In the current study, the number of bacterial 16S rRNA gene copies increased in the mesophilic and thermophilic environments compared to the psychrophilic environment (Table 1), suggesting that elevated temperature might facilitate the biotic degradation of cephapirin via the secretion of β -lactamases by an increased population of manure microbes.

On day 1, cephapirin concentration was lower in manure exposed to initial alkaline pH shock (pH 9 and 12) compared to neutral or acid pH shock (pH 7 or 5; Figure 1B). Gilbertson et al. (1990) demonstrated that the hydrolysis of ceftiofur was dramatically increased when pH was changed from pH 5 to 9. Ivaska and Nordström (1983) found that cephapirin exhibited a stable peak on differential pulse polarography at pH 7 to 8.5, implying that the chemical structure of cephapirin is more stable in a slightly alkaline condition.

In the current study, none of the animals had received any antibiotic treatment for at least 280 days before the cephapirin administration, but all three of the ARGs quantified (*cfxA*, *mefA*, and *tet(W)*) were detected in all of the manure slurry samples. Muurinen et al. (2017) detected 8 common ARGs and mobile genetic elements in stored manure as well as in soil (including fertilized and unfertilized) and tile drainage water collected from 2 dairy farms, which supports the co-occurrence of ARGs in various farm environments. It is likely that the detected ARGs could be transferred to the cows from the dairy farm environment or from other cows in the dairy herd, and therefore, various modes of ARG transmission deserve further attention and research.

Tetracycline resistance genes are the common ARGs observed in dairy manure (Sun et al., 2016; Zhou et al., 2016). In the current study, the average log base 10 transformed relative abundances for *tet(W)*, *mefA*, and *cfxA* cross all the temperature and pH treatments were -1.8, -2.2, and -4.3, respectively. A greater abundance of the *tet(W)* gene was detected than of the *cfxA* or *mefA* gene. The *cfxA* gene encodes class A β -lactamase, a gene that codes for resistance specific to cephalosporin antibiotics (García et al., 2008). In the current study, the *cfxA* gene had a low abundance compared to the abundance of *tet(W)* and *mefA* genes. This could be due to the rapid dissipation of cephapirin during manure incubation leading to low or no selective pressure among manure microbes.

No significant temperature or initial pH effects on the relative abundance of *cfxA* and *tet(W)* were observed in dairy manure slurry (Table 1). Sun et al. (2016) reported that there was no significant temperature effect on *tet(W)* abundance during anaerobic digestion of dairy manure. Huang et al. (2019) demonstrated that high temperature cannot always remove ARGs in manure anaerobic digestion unless antibiotic resistant bacteria or gene transfer elements are more efficiently decreased by raising temperature. Therefore, we can conclude that manure bacteria that encoded *cfxA* or *tet(W)* genes were not sensitive to incubation temperature or initial pH shock in dairy manure slurries.

Compared to the mesophilic condition, the *mefA* gene abundance decreased in psychrophilic and thermophilic environments ($P < 0.0001$; Table 1) probably due to changes in bacteria community (Qian et al., 2016; Sun et al., 2016). Miller et al. (2016) indicated that high temperature could remove some ARGs carried by bacterial hosts that were not thermotolerant. The optimum growth temperature for most bacterial species is around 37°C (Zhu, 2000). Therefore, the psychrophilic and thermophilic environments might inhibit the growth rate of bacteria that encoded *mefA* gene, leading to the decrease of *mefA* abundance.

The strong alkaline treatment (pH 12) resulted in 0.9 log base 10 fold reduction of *mefA* abundance as compared to the other pH treatments ($P < 0.0001$; Table 1). Although there was no pH effect on total bacterial 16S rRNA gene copies (Table 1), it is possible that the microbial community structure shifted due to pH stress. Lin et al. (2013) reported that bacteria *Clostridium alkalicellum* and *Corynebacterium humireducens* were enriched at pH 10 in swine manure, while the abundance of *Butyrivibrio* *sp.* was decreased. In the current study, the strong alkaline condition might have changed the bacterial community to have less bacteria carrying *mefA* gene, which led to the reduction in *mefA* abundance.

In conclusion, the thermophilic temperatures and strong alkaline treatments can facilitate rapid dissipation of cephalosporins. Both psychrophilic and thermophilic temperatures as well as strong alkaline treatments can decrease the abundance of *mefA* ARG. However, cephalosporins were rapidly and completely removed from manure slurries regardless of heat and pH treatment and thus, likely, no additional treatments are needed for its degradation. On the other hand, heat and pH influenced the abundance of *mefA* gene. This indicates that there could be treatment strategies (such as change in temperature or initial pH adjustment) developed to aid the removal of *mefA*. For instance, adding hydrated lime into manure slurries could be an effective method to reduce antibiotic resistance, since it will generate heat and increase pH. Heinonen-Tanski et al. (2004) reported that a dose of 10 g/L of hydrated lime with good stirring could destroy coliform bacteria to concentrations below 10 CFU/g in diluted cattle slurries. Jamal et al. (2011) indicated that adjusting pH to 12 or higher can halt or retard the microbial reactions leading to decreased odor production and vector attraction. Meanwhile, high pH can also reduce the availability of heavy metals, enhancing agricultural benefits and lowering environmental risks (Wong and Selvam, 2006). However, the role of elevated manure temperature and pH to nitrogen losses via volatilization of ammonia should be mentioned. The significant volatilization of ammonia during manure liming is a safety concern for animals and workers, and the loss of N also represents a significant loss of value for the manure as a fertilizer. Therefore, large stores of manure slurry should never be treated with lime due to a safety concern and an economic loss. Rather, liming of manure might be conducted within small loads of manure being hauled to a land application site in a tank wagon. One potential limitation of this study is that manure samples that had been treated to adjust pH were not neutralized before freezing. Without neutralizing samples at the completion of the incubation time, pH might impact the samples during storage and later

analysis. Although immediate storage at -20°C would stop almost all microbial growth and chemical properties of fresh manure (Pan et al., 2019), it may be worth highlighting that -80°C instead of -20°C is the best storage temperature for microbial community analyses, and pH may have impacted the degradation of cephalosporins. Overall, our results suggest increasing temperature or pH during storage of dairy manure slurries could be used together with other on-farm practices that are tailored to reduce the transfer of ARGs from manure to the environment following land application.

Acknowledgements

This study was supported by NIFA Competitive Grant no. 2014-05280 from the USDA National Institute of Food and Agriculture.

Reference

- Alcaine, S. D., S. S. Sukhnanand, L. D. Warnick, W.-L. Su, P. McGann, P. McDonough, and M. Wiedmann. 2005. Ceftiofur-resistant *Salmonella* strains isolated from dairy farms represent multiple widely distributed subtypes that evolved by independent horizontal gene transfer. *Antimicrob. Agents Chemother.* 49(10):4061-4067.
- Aminov, R., N. Garrigues-Jeanjean, and R. Mackie. 2001. Molecular ecology of tetracycline resistance: development and validation of primers for detection of tetracycline resistance genes encoding ribosomal protection proteins. *Appl. Environ. Microbiol.* 67(1):22-32.
- Cha, J., S. Yang, and K. Carlson. 2006. Trace determination of β -lactam antibiotics in surface water and urban wastewater using liquid chromatography combined with electrospray tandem mass spectrometry. *J. Chromatogr. A* 1115(1-2):46-57.
- Fahrenfeld, N., K. Knowlton, L. A. Krometis, W. C. Hession, K. Xia, E. Lipscomb, K. Libuit, B. L. Green, and A. Pruden. 2014. Effect of manure application on abundance of antibiotic

222 resistance genes and their attenuation rates in soil: field-scale mass balance approach. *Environ.*
 223 *Sci. Technol.* 48(5):2643-2650.

224 García, N., G. Gutiérrez, M. Lorenzo, J. E. García, S. Píriz, and A. Quesada. 2008. Genetic
 225 determinants for *cfxA* expression in *Bacteroides* strains isolated from human infections. *J.*
 226 *Antimicrob. Chemother.* 62(5):942-947.

227 Garfí, M., J. Martí-Herrero, A. Garwood, and I. Ferrer. 2016. Household anaerobic digesters for
 228 biogas production in Latin America: A review. *Renew. Sust. Energ. Rev.* 60:599-614.

229 Gilbertson, T. J., R. E. Hornish, P. S. Jaglan, K. T. Koshy, J. L. Nappier, G. L. Stahl, A. R.
 230 Cazars, J. M. Nappier, and M. F. Kubicek. 1990. Environmental fate of ceftiofur sodium, a
 231 cephalosporin antibiotic. Role of animal excreta in its decomposition. *J. Agric. Food Chem.*
 232 38(3):890-894.

233 Gou, M., H.-W. Hu, Y.-J. Zhang, J.-T. Wang, H. Hayden, Y.-Q. Tang, and J.-Z. He. 2018.
 234 Aerobic composting reduces antibiotic resistance genes in cattle manure and the resistome
 235 dissemination in agricultural soils. *Sci. Total Environ.* 612:1300-1310.

236 Heinonen-Tanski, H., S. Antola, and K. Weppling. 2004. Hydrated lime and Velox rapidly
 237 reduce enteric microorganisms in manure. *Sustainable Organic Waste Management for*
 238 *Environmental Protection and Food Safety. RAMRAN 2004, FAO and CSIC*, 2:33-36.

239 Hu, H. W., J. T. Wang, J. Li, J. J. Li, Y. B. Ma, D. Chen, and J. Z. He. 2016. Field-based
 240 evidence for copper contamination induced changes of antibiotic resistance in agricultural soils.
 241 *Environ. Microbiol.* 18(11):3896-3909.

242 Huang, X., J. Zheng, S. Tian, C. Liu, L. Liu, L. Wei, H. Fan, T. Zhang, L. Wang, and G. Zhu.
 243 2019. Higher Temperatures Do Not Always Achieve Better Antibiotic Resistance Gene Removal
 244 in Anaerobic Digestion of Swine Manure. *Appl. Environ. Microbiol.* 85(7):e02878-02818.

245 Ivaska, A. and F. Nordström. 1983. Determination of some cephalosporins by differential pulse
 246 polarography and linear scan voltammetry. *Anal. Chim. Acta* 146:87-95.
 247 Jamal, A., N. Norieh, and M. Farzadkia. 2011. Comparison of aerobic and lime stabilization
 248 methods for evaluation of sewage sludge reuse. *J. Environ. Sci. Technol.* 4(2):182-190.
 249 Kemper, N. 2008. Veterinary antibiotics in the aquatic and terrestrial environment. *Ecol. Indic.*
 250 8(1):1-13.
 251 Knapp, C. W., C. A. Engemann, M. L. Hanson, P. L. Keen, K. J. Hall, and D. W. Graham. 2008.
 252 Indirect evidence of transposon-mediated selection of antibiotic resistance genes in aquatic
 253 systems at low-level oxytetracycline exposures. *Environ. Sci. Technol.* 42(14):5348-5353.
 254 Li, R., J. A. Jay, and M. K. Stenstrom. 2019. Fate of antibiotic resistance genes and antibiotic-
 255 resistant bacteria in water resource recovery facilities. *Water Environ. Res.* 91(1):5-20.
 256 Lin, L., C. Wan, X. Liu, D.-J. Lee, Z. Lei, Y. Zhang, and J. H. Tay. 2013. Effect of initial pH on
 257 mesophilic hydrolysis and acidification of swine manure. *Bioresour. Technol.* 136:302-308.
 258 Looft, T., T. A. Johnson, H. K. Allen, D. O. Bayles, D. P. Alt, R. D. Stedtfeld, W. J. Sul, T. M.
 259 Stedtfeld, B. Chai, and J. R. Cole. 2012. In-feed antibiotic effects on the swine intestinal
 260 microbiome. *Proc. Natl. Acad. Sci.* 109(5):1691-1696.
 261 Martínez, J. L. 2008. Antibiotics and antibiotic resistance genes in natural environments. *Science*
 262 321(5887):365-367.
 263 Miller, J. H., J. T. Novak, W. R. Knocke, and A. Pruden. 2016. Survival of antibiotic resistant
 264 bacteria and horizontal gene transfer control antibiotic resistance gene content in anaerobic
 265 digesters. *Front. Microbiol.* 7:263.

266 Muurinen, J., R. Stedtfeld, A. Karkman, K. Pärnänen, J. Tiedje, and M. Virta. 2017. Influence of
 267 manure application on the environmental resistome under Finnish agricultural practice with
 268 restricted antibiotic use. *Environ. Sci. Technol.* 51(11):5989-5999.

269 Pan, J., J. Fadel, R. Zhang, H. El-Mashad, Y. Ying, and T. Rumsey. 2009. Evaluation of sample
 270 preservation methods for poultry manure. *Poult. Sci.* 88(8):1528-1535.

271 Pandyaswargo, A. H. and D. G. J. Premakumara. 2014. Financial sustainability of modern
 272 composting: the economically optimal scale for municipal waste composting plant in developing
 273 Asia. *Int. J. Recycl. Org. Waste Agric.* 3(3):4.

274 Pinheiro, J., D. Bates, S. DebRoy, D. Sarkar, and R. C. Team. 2018. nlme: Linear and nonlinear
 275 mixed effects models. R package version version 3.1-137.

276 Qian, X., W. Sun, J. Gu, X.-J. Wang, Y.-J. Zhang, M.-L. Duan, H.-C. Li, and R.-R. Zhang. 2016.
 277 Reducing antibiotic resistance genes, integrons, and pathogens in dairy manure by continuous
 278 thermophilic composting. *Bioresour. Technol.* 220:425-432.

279 R development Core Team. 2015. R: A language and environment for statistical computing.
 280 <http://www.R-project.org>.

281 Ray, P., C. Chen, K. F. Knowlton, A. Pruden, and K. Xia. 2017. Fate and effect of antibiotics in
 282 beef and dairy manure during static and turned composting. *J. Environ. Qual.* 46(1):45-54.

283 Ray, P., K. F. Knowlton, C. Shang, and K. Xia. 2014. Development and validation of a UPLC-
 284 MS/MS method to monitor cephalixin excretion in dairy cows following intramammary infusion.
 285 *PloS one* 9(11):e112343.

286 Sandegren, L. 2014. Selection of antibiotic resistance at very low antibiotic concentrations. *Ups.*
 287 *J. Med. Sci.* 119(2):103-107.

288 S3ki, J., S. M. Gonzalez, E. Urb3n, E. Nagy, and J. A. Ayala. 2011. Molecular analysis of the
 289 effector mechanisms of cefoxitin resistance among Bacteroides strains. J. Antimicrob.
 290 Chemother. 66(11):2492-2500.

291 Sun, W., X. Qian, J. Gu, X.-J. Wang, and M.-L. Duan. 2016. Mechanism and effect of
 292 temperature on variations in antibiotic resistance genes during anaerobic digestion of dairy
 293 manure. Sci. Rep. 6:30237.

294 Suzuki, M. T., L. T. Taylor, and E. F. DeLong. 2000. Quantitative analysis of small-subunit
 295 rRNA genes in mixed microbial populations via 5'-nuclease assays. Appl. Environ. Microbiol.
 296 66(11):4605-4614.

297 Tello, A., B. Austin, and T. C. Telfer. 2012. Selective pressure of antibiotic pollution on bacteria
 298 of importance to public health. Environ. Health Perspect. 120(8):1100-1106.

299 USFDA. 2018. 2017 summary report on antimicrobials sold or distributed for use in food-
 300 producing animals.
 301 [https://www.fda.gov/downloads/ForIndustry/UserFees/AnimalDrugUserFeeActADUFA/UCM62](https://www.fda.gov/downloads/ForIndustry/UserFees/AnimalDrugUserFeeActADUFA/UCM628538.pdf)
 302 [8538.pdf](https://www.fda.gov/downloads/ForIndustry/UserFees/AnimalDrugUserFeeActADUFA/UCM628538.pdf).

303 Wagner, R. D., S. J. Johnson, C. E. Cerniglia, and B. D. Erickson. 2011. Bovine intestinal
 304 bacteria inactivate and degrade ceftiofur and ceftriaxone with multiple β -lactamases. Antimicrob.
 305 Agents Chemother. 55(11):4990-4998.

306 Wong, J.W.C. and A. Selvam, 2006. Speciation of heavy metals during co-composting of sewage
 307 sludge with lime. Chemosphere, 63: 980-986.

308 Youngquist, C. P., S. M. Mitchell, and C. G. Cogger. 2016. Fate of antibiotics and antibiotic
 309 resistance during digestion and composting: a review. J. Environ. Qual. 45(2):537-545.

310 Zhou, B., C. Wang, Q. Zhao, Y. Wang, M. Huo, J. Wang, and S. Wang. 2016. Prevalence and
311 dissemination of antibiotic resistance genes and coselection of heavy metals in Chinese dairy
312 farms. *J. Hazard. Mater.* 320:10-17.

313 Zhu, J. 2000. A review of microbiology in swine manure odor control. *Agric., Ecosyst. Environ.*
314 78(2):93-106.




Figure 1. Cephapirin concentrations in dairy manure slurries exposed to various temperatures (A) or initial pH shocks (B) on day 1 following treatments ($n = 4$). Bars denoted by a different letter indicate significant differences between treatments ($P < 0.05$).

Table 1. The effect of temperature or initial pH on the abundance of 16S rRNA gene and relative abundance of antibiotic resistance genes in dairy manure slurries¹

Treatment	Bacterial 16S rRNA	<i>cfxA</i>	<i>mefA</i>	<i>tet(W)</i>
Temperature, °C				
10	8.28 ^a	-4.03	-2.40 ^a	-1.68
35	8.77 ^b	-4.19	-1.70 ^b	-1.80
55	8.64 ^b	-4.22	-2.69 ^a	-1.65
SEM ²	0.07	0.08	0.22	0.06
<i>P</i> value for temperature	< 0.0001	0.12	< 0.0001	0.19
Initial pH				
5	8.72	-4.30	-1.97 ^b	-1.94
7	8.75	-4.51	-1.92 ^b	-1.89
9	8.51	-4.32	-1.84 ^b	-1.58
12	8.69	-4.82	-2.81 ^a	-1.99
SEM	0.16	0.22	0.10	0.17
<i>P</i> value for initial pH	0.17	0.3	< 0.0001	0.25

¹Bacterial 16S rRNA gene was expressed as gene copies per g dry manure slurry. *cfxA* is a beta-lactam antibiotic resistance gene; *mefA* is a macrolide antibiotic resistance gene; and *tet(W)* is a tetracycline antibiotic resistance gene. Antibiotic resistance genes were expressed as a proportion to bacterial 16S rRNA gene. Data was normalized using log base 10 transformation. The Tukey's post hoc test was used to test differences among treatments. Different superscripts in the same column within temperature or pH group indicate significantly different means ($P < 0.05$). n = 4.

²Standard error of mean